

# Pseudozero Set of Multivariate Polynomials

Stef Graillat

Laboratoire LP2A, Université de Perpignan Via Domitia  
52, avenue Paul Alduy, F-66860 Perpignan cedex, France  
graillat@univ-perp.fr  
<http://gala.univ-perp.fr/~graillat>

**Abstract.** The pseudozero set of a system  $P$  of polynomials in  $n$  variables is the subset of  $\mathbb{C}^n$  consisting of the union of the zeros of all polynomial systems  $Q$  that are near to  $P$  in a suitable sense. This concept arises naturally in Scientific Computing where data often have a limited accuracy. When the polynomials of the system are polynomials with complex coefficients, the pseudozero set has already been studied. In this paper, we focus on the case where the polynomials of the system have real coefficients and such that all the polynomials in all the perturbed polynomial systems have real coefficients as well. We provide an explicit definition to compute this pseudozero set. At last, we analyze different methods to visualize this set.

## 1 Introduction and notation

### 1.1 Summary

Polynomials appear in almost all areas in scientific computing and engineering as it is shown in the Computer Algebra Handbook [6] and in [4]. Most of the applications need to solve equations involving polynomials and systems of polynomials, often in many variables. The relationships between industrial applications and polynomial systems solving were studied by the European Community Project FRISCO. The report may be found at <http://www.nag.co.uk/projects/FRISCO.html>. They gave a list of the major fields where polynomial systems are used: Computer Aided Design and Modeling, Mechanical Systems Design, Signal Processing and Filter Design, Civil Engineering, Robotics, Simulation. The wide range of use of polynomial systems needs to have fast and reliable methods to solve them. Roughly speaking, there are two general approaches: symbolic and numeric. The symbolic approach is based either on the theory of Gröbner basis or on the theory of resultants. For the numeric approach, it is the use of iterative methods like Newton's method or homotopy continuation methods. Recently, hybrid methods, combining both symbolic and numeric methods, began to appear (see the chapter called "Hybrid Methods" by Kaltofen in [6, p.112-128]).

In practice, from situations arising in science or engineering, the data are known only to a limited accuracy. From a polynomial point of view, this only

means that the coefficients of the polynomials are known only to within a certain tolerance. Then it is important to obtain informations about the variation of the zeros of the polynomial or of the polynomial system in the presence of uncertainty on the coefficients. Analytical sensitivity analysis introduces a condition number that bounds the magnitudes of the (first order) changes of the roots with respect to the coefficient perturbations. Numerous results in this direction are available, see for example Gautschi [5] or Wilkinson [26]. Representing coefficient uncertainty with intervals and computing with interval arithmetic yield over-sets that enclose (sometimes pessimistically) the perturbed roots. Continuous sensitivity analysis, introduced by Ostrowski [19], considers the uncertainty of the coefficients as a continuity problem. The most powerful tool of this last type of methods seems to be the *pseudozero set* of a polynomial we focus hereafter. Roughly speaking, this is the set of roots of polynomials that are near to a given polynomial.

The pseudozero set was first introduced by Mosier [18] in 1986. He studied this set considering perturbations bounded with the  $\infty$ -norm. Trefethen and Toh [25] studied pseudozero set for perturbations bounded with the 2-norm. They also compared the pseudozero set of a given polynomial with the pseudospectra of the associated companion matrix. These results are summarized in Chatelin and Frayssé's book on finite precision [2]. More recently, Zhang [27] compared pseudozero set with respect to the choice of the polynomial basis (power, Taylor, Chebyshev, Bernstein). At last, recently, Stetter gave a general framework for working with inexact polynomials in his book [24] (based on previous papers [21–23]). The notion of root sets was introduced by Hinrichsen and Kelb [12]. It is a particular case of the spectral value sets of the companion matrix using structured perturbations. It corresponds exactly to the notion of pseudozero set but from a different viewpoint. Such a set was studied in particular by Hinrichsen and Kelb [12], Karow [16] and Hinrichsen and Pritchard [13].

Nevertheless, few applications of pseudozero set have been given in these previous publications, except when Bini and Fiorentino provided a multiprecision algorithm to compute polynomial root using pseudozero set [1]. Indeed, they need to know if an approximate root is a root of a nearby polynomial. Pseudozero set is the natural way to answer this question. More recently, Graillat and Langlois [7–10] gave some applications of pseudozero set in Computer Algebra and in Control Theory. They provide in these articles an algorithm to test the approximate primality of two univariate polynomials. They also propose an algorithm to compute the stability radius of an univariate polynomial.

The major part of the papers cited above consider only the univariate case. The multivariate case seems to have received few attention. It has only been studied by Stetter in [22, 24], by Hoffman, Madden and Zhang in [14] and Corless, Kai and Watt in [3]. Furthermore, the multivariate case has only been dealt with polynomials with complex coefficients. In this paper, we consider systems where polynomials have real coefficients and such that all the polynomials in all the perturbed polynomial systems have real coefficients as well. We provide a simple

criterion to compute the pseudozero set and study different methods to visualize it.

The rest of the paper is organized as follows. In the rest of this Section, we introduce some notations and well-known results on basic linear algebra and computer algebra. In Section 2, we recall results on complex pseudozero set. In Section 3, we study real pseudozero set and establish a computable criterion for this pseudozero set. In Section 4, we present different methods to visualize the pseudozero set.

## 1.2 Notation

We recall the notations used in Stetter [24]. A *monomial* in the  $n$  variables  $z_1, \dots, z_n$  is the power product

$$z^j := z_1^{j_1} \cdots z_n^{j_n}, \quad \text{with } j = (j_1, \dots, j_n) \in \mathbb{N}^n;$$

$j$  is the *exponent* and  $|j| := \sum_{\sigma=1}^n j_\sigma$  the *degree* of the monomial  $z^j$ .

**Definition 1.** A complex (real) polynomial in  $n$  variables is a finite linear combination of monomials in  $n$  variables with coefficients from  $\mathbb{C}$  (from  $\mathbb{R}$ ),

$$p(z) = p(z_1, \dots, z_n) = \sum_{(j_1, \dots, j_n) \in J} a_{j_1 \dots j_n} z_1^{j_1} \cdots z_n^{j_n} = \sum_{j \in J} a_j z^j.$$

The set  $J \subset \mathbb{N}^n$  which contains the exponents of those monomials which are present in the polynomial  $p$  is called the *support* of  $p$ . The total *degree* of  $p$  is defined to be the number  $\deg(p) := \max_{j \in J} |j|$ . The set of all complex (real) polynomials in  $n$  variables will be denoted by  $\mathcal{P}^n(\mathbb{C})$  (by  $\mathcal{P}^n(\mathbb{R})$ ). When the coefficient domain is evident or is not important, the notation  $\mathcal{P}^n$  will be used. The notation  $\mathcal{P}_d^n \subset \mathcal{P}^n$  stands for the set of polynomials in  $n$  variables of total degree  $\leq d$ . As we will often manipulate polynomials with linear operations, we will widely employ the notations of linear algebra. We will generally collect the coefficients of a polynomial into a vector  $a = (\dots, a_j, \dots, j \in J)^T$  and its monomials into a vector  $\mathbf{z} = (\dots, z^j, \dots, j \in J)^T$ .

Let  $p = \sum_{j \in J} a_j z^j \in \mathcal{P}^n(\mathbb{K})$  with  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$  be a polynomial in  $n$  variables and  $J$  be its support. We denote by  $|J|$  the number of elements of  $J$ . If  $|J| = M$  and let  $\|\cdot\|$  be a norm on  $\mathbb{K}^M$ , we denote by  $\|p\|$  the norm of the vector  $a = (\dots, a_j, \dots, j \in J)$ , namely,

$$\|p\| := \|(\dots, a_j, \dots, j \in J)^T\|.$$

Given such an  $\varepsilon > 0$ , the  $\varepsilon$ -neighborhood  $N_\varepsilon(p)$  of the polynomial  $p \in \mathcal{P}^n(\mathbb{K})$  is the set of all polynomials of  $\mathcal{P}^n(\mathbb{K})$ , close enough to  $p$ , that is to say, the set of polynomials  $\tilde{p} = \sum_{j \in \tilde{J}} \tilde{a}_j z^j \in \mathcal{P}^n(\mathbb{K})$  with support  $\tilde{J} \subset J$  and  $\|\tilde{p} - p\| \leq \varepsilon$ .

Given a norm  $\|\cdot\|$  on  $\mathbb{K}^N$  with  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ , we define its *dual norm* (denoted by  $\|\cdot\|_*$ ) by

$$\|x\|_* := \sup_{y \neq 0} \frac{|y^T x|}{\|y\|} = \sup_{\|y\|=1} |y^T x|.$$

Table 1 represents the most common norms on  $\mathbb{K}^N$  and their respective dual norms. Given a vector  $x \in \mathbb{K}^N$ , there exists a vector  $y \in \mathbb{K}^N$  with  $\|y\| = 1$

Norms	Dual norms
$\ x\ _1 := \sum_j  x_j $	$\ x\ _1^* = \max_j  x_j  = \ x\ _\infty$
$\ x\ _2 := (\sum_j  x_j ^2)^{1/2}$	$\ x\ _2^* = (\sum_j  x_j ^2)^{1/2} = \ x\ _2$
$\ x\ _\infty := \max_j  x_j $	$\ x\ _\infty^* = \sum_j  x_j  = \ x\ _1$

**Table 1.** Dual norms for most common norms on  $\mathbb{K}^N$

satisfying  $x^T y = \|x\|_*$  (see [11, p.107] or [15, p. 278]). The vector  $y$  is called the *dual vector* of  $x$ .

**Definition 2.** A value  $z \in \mathbb{K}^n$  is a  $\varepsilon$ -pseudozero of a polynomial  $p \in \mathcal{P}^n$  if it is a zero of some polynomial  $\tilde{p}$  in  $N_\varepsilon(p)$ .

**Definition 3.** The  $\varepsilon$ -pseudozero set of a polynomial  $p \in \mathcal{P}^n$  (denoted by  $Z_\varepsilon(p)$ ) is the set of all the  $\varepsilon$ -pseudozeros,

$$Z_\varepsilon(p) := \{z \in \mathbb{K}^n : \exists \tilde{p} \in N_\varepsilon(p), \tilde{p}(z) = 0\}.$$

Three important issues arise from these definitions.

- For  $p$  with real coefficients  $a_j$ , it must be specified whether  $N_\varepsilon(p)$  is restricted to real polynomials and not. Indeed, it seems natural for a real polynomial to be perturbed by real polynomials.
- One may only be interested in real or complex pseudozero set.
- The pseudozero set  $Z_\varepsilon(p)$  cannot be computed directly because it is the union of the zeros of an infinite number of polynomials.

We can extend those definitions to a *system* of polynomials

$$P = \{p_1, \dots, p_k\}, \quad k \in \mathbb{N}.$$

We will often consider this system as a vectors of polynomials

$$P(z) = \begin{pmatrix} p_1(z) \\ \vdots \\ p_k(z) \end{pmatrix}.$$

Given an  $\varepsilon > 0$  and a *system* of polynomials  $P = \{p_1, \dots, p_k\}$ ,  $k \in \mathbb{N}$ , the  $\varepsilon$ -neighborhood  $N_\varepsilon(P)$  is the set of systems of polynomials  $\tilde{P} = \{\tilde{p}_1, \dots, \tilde{p}_k\}$  close enough to  $P$ , that is to say with  $\tilde{p}_j \in N_\varepsilon(p_j)$  for  $j = 1, \dots, k$ .

**Definition 4.** A value  $z \in \mathbb{K}^n$  is a  $\varepsilon$ -pseudozero of a polynomial system  $P$  if it is a zero of a system of polynomials  $\tilde{P}$  in  $N_\varepsilon(P)$ .

**Definition 5.** The  $\varepsilon$ -pseudozero set of a system of polynomials  $P$  (denoted by  $Z_\varepsilon(P)$ ) is the set of all the  $\varepsilon$ -pseudozeros,

$$Z_\varepsilon(P) := \{z \in \mathbb{K}^n : \exists \tilde{P} \in N_\varepsilon(P), \tilde{P}(z) = 0\}.$$

## 2 Pseudozero set of complex multivariate polynomials

Theorem 1 below provides a computable counterpart of the pseudozero set.

**Theorem 1 (Stetter [24]).** *The complex  $\varepsilon$ -pseudozero set of  $p = \sum_{j \in J} a_j z^j \in \mathcal{P}^n(\mathbb{C})$  verifies*

$$Z_\varepsilon(p) = \left\{ z \in \mathbb{C}^n : g(z) := \frac{|p(z)|}{\|\mathbf{z}\|_*} \leq \varepsilon \right\},$$

where  $\mathbf{z} := (\dots, |z|^j, \dots, j \in J)^T$ .

For completeness of the paper, we recall the proof.

*Proof.* If  $z \in Z_\varepsilon(p)$  then there exists  $\tilde{p} \in \mathcal{P}^n$  such that  $\tilde{p}(z) = 0$  and  $\|p - \tilde{p}\| \leq \varepsilon$ . From the generalized Hölder's inequality  $|x^T y| \leq \|x\| \|y\|_*$ , we get

$$|p(z)| = |p(z) - \tilde{p}(z)| = \left| \sum_{j \in J} (p_j - \tilde{p}_j) z^j \right| \leq \|p - \tilde{p}\| \|\mathbf{z}\|_*.$$

It follows that  $|p(z)| \leq \varepsilon \|\mathbf{z}\|_*$ .

Conversely, let  $u \in \mathbb{C}$  be such that  $|p(u)| \leq \varepsilon \|\mathbf{u}\|$  where  $\mathbf{u} := (\dots, |u|^j, \dots, j \in J)$ . The dual vector  $d$  of  $\mathbf{u}$  verifies  $d^T \mathbf{u} = \|\mathbf{u}\|_*$  and  $\|d\| = 1$ . Let us introduce the polynomials  $r$  and  $p_u$  defined by

$$r(z) = \sum_{k=0}^n r_k z^k \text{ with } r_k = d_k,$$

$$p_u(z) = p(z) - \frac{p(u)}{r(u)} r(z).$$

This polynomial  $p_u$  is (with respect to the norm  $\|\cdot\|$ ) the nearest polynomial of  $p$  with  $u$  as a root. It is clear that  $r(u) = d^T \mathbf{u} = \|\mathbf{u}\|_*$ . So we have

$$\|p - p_u\| = \frac{|p(u)|}{|r(u)|} \|r\| \leq \varepsilon \|d\|.$$

As  $\|d\| = 1$ , we get

$$\|p - p_u\| \leq \varepsilon.$$

And since  $p_u(u) = 0$ ,  $u$  belongs to  $Z_\varepsilon(p)$ . □

This theorem can be immediately extended to systems of polynomials.

**Corollary 1 (Stetter [22]).** *The complex  $\varepsilon$ -pseudozero set of  $P = \{p_1, \dots, p_k\}$ ,  $k \in \mathbb{N}$  verifies*

$$Z_\varepsilon(P) = \left\{ z \in \mathbb{C}^n : \frac{|p_l(z)|}{\|\mathbf{z}_l\|_*} \leq \varepsilon \text{ for } l = 1, \dots, k \right\},$$

where  $\mathbf{z}_l := (\dots, |z|^j, \dots, j \in J_l)^T$ .

For the next theorem, we will restrict our attention to situations where  $P$  as well as all the systems in  $N_\varepsilon(P)$  are 0-dimensional, that is to say if the solutions of the system are non-empty and finite.

**Theorem 2 (Stetter [22]).** *Under the above assumptions, each system  $\tilde{P} \in N_\varepsilon(P)$  has the same number of zeros (counting multiplicities) in a fixed pseudozero set connected component of  $Z_\varepsilon(P)$ .*

*Proof.* This is the proof of [22, Thm. 3.5]. □

### 3 Pseudozero set of real multivariate polynomials

#### 3.1 Complex pseudozero set of real multivariate polynomials

A real  $\varepsilon$ -neighborhood of  $p$  is the set of all polynomials of  $\mathcal{P}^n(\mathbb{R})$ , close enough to  $p$ , that is to say,

$$N_\varepsilon^R(p) = \{\tilde{p} \in \mathcal{P}^n(\mathbb{R}) : \|p - \tilde{p}\| \leq \varepsilon\}.$$

Then the real  $\varepsilon$ -pseudozero set of  $p$  is defined to include all the zeros of the real  $\varepsilon$ -neighborhood of  $p$ . A definition of this set is

$$Z_\varepsilon^R(p) = \{z \in \mathbb{C}^n : \tilde{p}(z) = 0 \text{ for } \tilde{p} \in N_\varepsilon^R(p)\}.$$

For  $\varepsilon = 0$ , the pseudozero set  $Z_0^R(p)$  is the set of the roots of  $p$  we denote  $Z(p)$ .

Following Theorem 3 provides a computable counterpart of this definition. It is based on arguments developed by Hinrichsen and Kelb in [12]. We define for  $x, y \in \mathbb{R}^N$ ,

$$d(x, \mathbb{R}y) = \inf_{\alpha \in \mathbb{R}} \|x - \alpha y\|_*,$$

the distance of a point  $x \in \mathbb{R}^N$  from the linear subspace  $\mathbb{R}y = \{\alpha y, \alpha \in \mathbb{R}\}$ .

**Theorem 3.** *The real  $\varepsilon$ -pseudozero set of  $p = \sum_{j \in J} a_j z^j \in \mathcal{P}^n(\mathbb{R})$  verifies*

$$Z_\varepsilon^R(p) = Z(p) \cup \left\{ z \in \mathbb{C}^n \setminus Z(p) : h(z) := d(G_R(z), \mathbb{R}G_I(z)) \geq \frac{1}{\varepsilon} \right\},$$

where  $G_R(z)$  and  $G_I(z)$  are the real and imaginary parts of

$$G(z) = \frac{1}{p(z)} (\dots, z^j, \dots, j \in J)^T, \quad z \in \mathbb{C}^n \setminus Z(p).$$

*Proof.* Let  $z \in Z_\varepsilon^R(p)$ . If  $p(z) = 0$  then  $z \in Z(p)$  else there exists  $q \in N_\varepsilon^R(p)$  such that  $q(z) = 0$ . In this case, we have  $p(z) = p(z) - q(z) = (p - q)^T \underline{z}$ , where  $\underline{z} = (\dots, z^j, \dots, j \in J)^T$ . It follows that  $1 = (p - q)^T G(z)$ . Hence we have  $1 = (p - q)^T G_R(z) + i(p - q)^T G_I(z)$  and so

$$\begin{cases} (p - q)^T G_R(z) = 1, \\ (p - q)^T G_I(z) = 0. \end{cases}$$

As a consequence, we have  $\|p - q\| \|G_R(u) - \alpha G_I(u)\|_* \geq 1$ , for all  $\alpha \in \mathbb{R}$ . We conclude that

$$d(G_R(u), \mathbb{R}G_I(u)) \geq \frac{1}{\|p - q\|} \geq \frac{1}{\varepsilon}.$$

Conversely, let  $z \in Z(p) \cup \{z \in \mathbb{C}^n \setminus Z(p) : d(G_R(z), \mathbb{R}G_I(z)) \geq \frac{1}{\varepsilon}\}$ . If  $z$  belongs to  $Z(p)$  then it belongs to  $Z_\varepsilon^R(p)$ . Otherwise  $z$  satisfies  $d(G_R(z), \mathbb{R}G_I(z)) \geq 1/\varepsilon$ . From a duality theorem (see [17, p.119]), there exists a vector  $u \in \mathbb{R}^N$  with  $\|u\| = 1$  satisfying

$$u^T G_R(z) = d(G_R(z), \mathbb{R}G_I(z)) \quad \text{and} \quad u^T G_I(z) = 0.$$

Let us consider the real polynomial

$$q = p - \frac{u}{d(G_R(z), \mathbb{R}G_I(z))}.$$

We have

$$q(z) = p(z) - \frac{u^T z}{d(G_R(z), \mathbb{R}G_I(z))} = p(z) - \frac{p(z)u^T G(z)}{d(G_R(z), \mathbb{R}G_I(z))} = 0.$$

Furthermore we have  $\|q - p\| = 1/d(G_R(z), \mathbb{R}G_I(z))$ , so that  $\|p - q\| \leq \varepsilon$ .  $\square$

To compute the real  $\varepsilon$ -pseudozero set  $Z_\varepsilon^R(p)$ , we only have to evaluate the distance  $d(G_R(z), \mathbb{R}G_I(z))$ . This quantity can be calculated easily for the 2-norm. Let us now denote the 2-norm  $\|\cdot\|_2$  and  $\langle \cdot, \cdot \rangle$  the corresponding inner product. In this case, we have

$$d(x, \mathbb{R}y) = \begin{cases} \sqrt{\|x\|_2^2 - \frac{\langle x, y \rangle^2}{\|y\|_2^2}} & \text{if } y \neq 0, \\ \|x\|_2 & \text{if } y = 0. \end{cases}$$

For the  $\infty$ -norm, it is shown in [16, Prop. 7.7.2] that

$$d(x, \mathbb{R}y) = \begin{cases} \min_{\substack{i=0:n \\ y_i \neq 0}} \|x - (x_i/y_i)y\|_1 & \text{if } y \neq 0, \\ \|x\|_1 & \text{if } y = 0. \end{cases}$$

For the other  $p$ -norm with  $p \neq 2, \infty$ , there is no easy computable formula to calculate  $d(x, \mathbb{R}y)$ .

This theorem can be immediately extended to systems of polynomials.

**Corollary 2.** *The real  $\varepsilon$ -pseudozero set of  $P = \{p_1, \dots, p_k\}$ ,  $k \in \mathbb{N}$  verifies*

$$Z_\varepsilon^R(P) = \bigcap_{l=1}^k \left( Z(p_l) \cup \left\{ z \in \mathbb{C}^n \setminus Z(p_l) : h_l(z) := d(G_R^l(z), \mathbb{R}G_I^l(z)) \geq \frac{1}{\varepsilon} \right\} \right),$$

where  $G_R^l(z)$  and  $G_I^l(z)$  are the real and imaginary parts of

$$G^l(z) = \frac{1}{p_l(z)} (\dots, z^j, \dots, j \in J_l)^T, \quad z \in \mathbb{C}^n \setminus Z(p_l).$$

As we have seen before, the real pseudozero set is closely related to the function  $d$ . This function can have a discontinuous behavior. It is the subject of the following lemma.

**Lemma 1 (Hinrichsen and Kelb [12]).** *The function*

$$d : \mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}_+, \quad (x, y) \mapsto d(x, \mathbb{R}y)$$

*is continuous at all pairs  $(x, y)$  with  $y \neq 0$  or  $x = 0$  and discontinuous at all pairs  $(x, 0) \in \mathbb{R}^{n+1} \times \mathbb{R}^{n+1}$ ,  $x \neq 0$ .*

This lemma states that a discontinuity problem arises when vector  $y$  vanishes. In our case, the discontinuity arises when  $G_I(z) = 0$  where  $G_I(z)$  is the imaginary part of

$$G(z) = \frac{1}{p(z)}(1, z, \dots, z^n)^T.$$

It follows that  $G_I$  vanishes for  $z \in \mathbb{R}$ , that is along the real axis. This explains why the `contour` and `meshc` functions of MATLAB may fail and give some bad results along the real axis. Of course, if none of the zeros of the polynomial is real, the real pseudozero set is correct because we do not evaluate the function  $G$  on the real axis.

### 3.2 Real pseudozero set of real multivariate polynomials

In the previous subsection, we were interested in the complex zeros of a real polynomial system. Sometimes, we can be interested only in the real zeros of a system. That is to say, given a polynomial  $p \in \mathcal{P}^n(\mathbb{R})$ , we are interested in  $Z_\varepsilon^R(p) \cap \mathbb{R}^n$ . The following result gives a formula to compute this set.

**Theorem 4.** *The intersection between the complex  $\varepsilon$ -pseudozero set of  $p = \sum_{j \in J} a_j z^j \in \mathcal{P}^n(\mathbb{C})$  and  $\mathbb{R}^n$  verifies*

$$Z_\varepsilon^R(p) \cap \mathbb{R}^n = \left\{ z \in \mathbb{R}^n : g(z) := \frac{|p(z)|}{\|\mathbf{z}\|_*} \leq \varepsilon \right\},$$

where  $\mathbf{z} := (\dots, |z|^j, \dots, j \in J)^T$ .

*Proof.* The proof is the same as Theorem 1. □

This theorem can be immediately extended to systems of polynomials.

**Corollary 3.** *The intersection between the complex  $\varepsilon$ -pseudozero set of  $P = \{p_1, \dots, p_k\}$ ,  $k \in \mathbb{N}$  and  $\mathbb{R}^n$  verifies*

$$Z_\varepsilon^R(P) \cap \mathbb{R}^n = \left\{ z \in \mathbb{R}^n : \frac{|p_l(z)|}{\|\mathbf{z}_l\|_*} \leq \varepsilon \text{ for } l = 1, \dots, k \right\},$$

where  $\mathbf{z}_l := (\dots, |z|^j, \dots, j \in J_l)^T$ .

## 4 Visualization of pseudozero sets

The descriptions of  $Z_\varepsilon(P)$  and  $Z_\varepsilon^R(P)$  given by Theorem 1 and Theorem 3 (and by Corollary 1 and Corollary 2) enable us to compute, plot and visualize pseudozero set of multivariate polynomials. The pseudozero set is a subset of  $\mathbb{C}^n$  which can only be seen by its projections on low dimensional spaces that is often  $\mathbb{C}$ .

We have written a MATLAB program to compute and visualize these projections (see Appendix ??). This program requires the Symbolic Math Toolbox (and the Extended Symbolic Math Toolbox) which is the MATLAB gateway to the kernel of MAPLE.

For a given  $v \in \mathbb{C}^n$ , let  $Z_\varepsilon(P, j, v)$  be the projection of  $Z_\varepsilon(P)$  onto the  $z_j$ -space around  $v$ . Then, it follows that for  $P = \{p_1, \dots, p_k\}$ ,

$$Z_\varepsilon(P, j, v) = \left\{ z \in \mathbb{C}^n : z_i = v_i \text{ for } i \neq j, \text{ and } \max_{l=1, \dots, k} \frac{|p_l(z)|}{\|\mathbf{z}_l\|_*} \leq \varepsilon \right\},$$

where  $\mathbf{z}_l := (\dots, |z|^j, \dots, j \in J_l)^T$ . One way for visualizing  $Z_\varepsilon(P, j, v)$  is to plot the values of the projection of

$$\text{ps}(z) := \log_{10} \left( \max_{l=1, \dots, k} \frac{|p_l(z)|}{\|\mathbf{z}_l\|_*} \right)$$

over a set of grid points around  $v$  in  $z_j$ -space. In the same way, we define for a given  $v \in \mathbb{C}^n$ ,  $Z_\varepsilon^R(P, j, v)$  by the projection of  $Z_\varepsilon^R(P)$  onto the  $z_j$ -space around  $v$ . Then, it follows that for  $P = \{p_1, \dots, p_k\}$ ,

$$Z_\varepsilon^R(P, j, v) = \left\{ z \in \mathbb{C}^n : z_i = v_i \text{ for } i \neq j, \text{ and } \max_{l=1, \dots, k} d(G_R^l(z), \mathbb{R}G_I^l(z))^{-1} \leq \varepsilon \right\},$$

where  $G_R^l(z)$  and  $G_I^l(z)$  are the real and imaginary parts of

$$G^l(z) = \frac{1}{p_l(z)} (\dots, z^j, \dots, j \in J_l)^T, \quad z \in \mathbb{C}^n \setminus Z(p).$$


One way for visualizing  $Z_\varepsilon^R(P, j, v)$  is still to plot the values of the projection of

$$\text{ps}^R(z) := \log_{10} \left( \max_{l=1, \dots, k} d(G_R^l(z), \mathbb{R}G_I^l(z))^{-1} \right)$$

over a set of grid points around  $v$  in  $z_j$ -space. We examine the following system from [14] (see Figure 1) using the 2-norm: two unit balls intersection at  $(2, 2)$ ,

$$P_1 = \begin{cases} p_1 = (z_1 - 1)^2 + (z_2 - 2)^2 - 1, \\ p_2 = (z_1 - 3)^2 + (z_2 - 2)^2 - 1. \end{cases}$$

We can be only interested in the real zeros of a polynomial systems. In this case, we can only draw  $\mathbb{R}^n \cap Z_\varepsilon^R(P)$ . This is what is done with the following



**Fig. 1.** Projections of the complex pseudozero set (on the left) and the real pseudozero set (on the right) of  $P_1$


example in Figure 2 still with the 2-norm,

$$P_2 = \begin{cases} p_1 = z_1^2 + z_2^2 - 1, \\ p_2 = 25z_1z_2 - 12. \end{cases}$$

In this Figure, we have computed the function

$$g(x, y) = \max_{l=1,2} \frac{p_l(x, y)}{\|\mathbf{z}_l\|_*},$$

where  $\mathbf{z}_l := (\dots, |x + iy|^j, \dots, j \in J_l)^T$ .



**Fig. 2.** Projection of the real pseudozero set of  $P_2$

Several issues appear when one wants to draw the real or complex pseudozero set. First, one has to choose a discretization that separates the roots. It is often a difficult task. For drawing the real pseudozero set, one needs to deal with function  $d$  that is discontinuous on the real axis.

The cost of our algorithms can be very high because it depends on the number of nodes of the grid. Nevertheless, we are not interested in providing cheap algorithms. We just want to provide tools that enable us to make a qualitative analysis of a polynomial.

## 5 Conclusion

Approximate polynomials are unavoidable in numerous application fields and in finite precision environment. Plotting pseudozero set can give qualitative and

sometimes quantitative interesting informations about the behavior of these approximate polynomials. We have shown that pseudozero set offers a powerful tool. They can be easily plotted using popular software as MATLAB. We hope that pseudozero set will be used as much as pseudospectra.

## References

1. Dario Andrea Bini and Giuseppe Fiorentino. Design, analysis, and implementation of a multiprecision polynomial rootfinder. *Numer. Algorithms*, 23(2-3):127–173, 2000.
2. Françoise Chaitin-Chatelin and Valérie Frayssé. *Lectures on finite precision computations*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1996.
3. Robert M. Corless, Hiroshi Kai, and Stephen M. Watt. Approximate computation of pseudovarieties. *SIGSAM Bull.*, 37(3):67–71, 2003.
4. Alicia Dickenstein and Ioannis Z. Emiris, editors. *Solving polynomial equations: Foundations, algorithms, and applications*, volume 14 of *Algorithms and Computation in Mathematics*. Springer-Verlag, Berlin, 2005.
5. Walter Gautschi. On the condition of algebraic equations. *Numer. Math.*, 21:405–424, 1973.
6. Johannes Grabmeier, Erich Kaltofen, and Volker Weispfenning, editors. *Computer algebra handbook*. Springer-Verlag, Berlin, 2003.
7. Stef Graillat and Philippe Langlois. Testing polynomial primality with pseudozeros. In *Proceedings of the Fifth Conference on Real Numbers and Computers*, pages 231–246, Lyon, France, September 2003.
8. Stef Graillat and Philippe Langlois. Computation of stability radius for polynomials. Preprint 31, Université de Perpignan, January 2004.
9. Stef Graillat and Philippe Langlois. More on pseudozeros for univariate polynomials. Preprint 32, Université de Perpignan, January 2004.
10. Stef Graillat and Philippe Langlois. Pseudozero set decides on polynomial stability. In *Proceedings of the Symposium on Mathematical Theory of Networks and Systems*, Leuven, Belgium, July 2004.
11. Nicholas J. Higham. *Accuracy and Stability of Numerical Algorithms*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, second edition, 2002.
12. D. Hinrichsen and B. Kelb. Spectral value sets: a graphical tool for robustness analysis. *Systems Control Lett.*, 21(2):127–136, 1993.
13. Diederich Hinrichsen and Anthony J. Pritchard. *Mathematical systems theory. I*, volume 48 of *Texts in Applied Mathematics*. Springer-Verlag, Berlin, 2005. Modelling, state space analysis, stability and robustness.
14. J. William Hoffman, James J. Madden, and Hong Zhang. Pseudozeros of multivariate polynomials. *Math. Comp.*, 72(242):975–1002 (electronic), 2003.
15. Roger A. Horn and Charles R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, 1990.
16. Michael Karow. *Geometry of spectral value sets*. PhD thesis, Universität Bremen, 2003.
17. David G. Luenberger. *Optimization by vector space methods*. John Wiley & Sons Inc., New York, 1969.
18. Ronald G. Mosier. Root neighborhoods of a polynomial. *Math. Comp.*, 47(175):265–273, 1986.

19. A. M. Ostrowski. *Solution of equations and systems of equations*. Second edition. Pure and Applied Mathematics, Vol. 9. Academic Press, New York, 1966.
20. Hans J. Stetter. Analysis of zero clusters in multivariate polynomial systems. In *ISSAC '96: Proceedings of the 1996 international symposium on Symbolic and algebraic computation*, pages 127–136. ACM Press, 1996.
21. Hans J. Stetter. The nearest polynomial with a given zero, and similar problems. *SIGSAM Bulletin (ACM Special Interest Group on Symbolic and Algebraic Manipulation)*, 33(4):2–4, décembre 1999.
22. Hans J. Stetter. Polynomials with coefficients of limited accuracy. In *Computer algebra in scientific computing—CASC'99 (Munich)*, pages 409–430. Springer, Berlin, 1999.
23. Hans J. Stetter. Condition analysis of overdetermined algebraic problems. In *Computer algebra in scientific computing (Samarkand, 2000)*, pages 345–365. Springer, Berlin, 2000.
24. Hans J. Stetter. *Numerical Polynomial Algebra*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2004.
25. Kim-Chuan Toh and Lloyd N. Trefethen. Pseudozeros of polynomials and pseudospectra of companion matrices. *Numer. Math.*, 68(3):403–425, 1994.
26. J. H. Wilkinson. *Rounding errors in algebraic processes*. Dover Publications Inc., New York, 1994.
27. Hong Zhang. Numerical condition of polynomials in different forms. *Electron. Trans. Numer. Anal.*, 12:66–87 (electronic), 2001.